arXiv:0808.1619v4 [math.AG] 9 Nov 2009

COUPLED PAINLEVÉ SYSTEMS WITH AFFINE WEYL GROUP SYMMETRY OF TYPES $A_7^{(2)}, A_5^{(2)}$ AND $D_4^{(3)}$

Abstract. We find a four-parameter family of coupled Painlevé VI systems in dimension four with affine Weyl group symmetry of type $A_7^{(2)}$. This is the first example which gave higher-order Painlevé equations of type $A_{2l+5}^{(2)}$. We then give an explicit description of a confluence process from this system to a 3-parameter family of coupled Painlevé V and III systems in dimension four with $W(A_5^{(2)})$-symmetry. For a degenerate system of $A_5^{(2)}$ system, we also find a two-parameter family of ordinary differential systems in dimension four with affine Weyl group symmetry of type $D_4^{(3)}$. This is the first example which gave higher-order Painlevé equations of type $D_4^{(3)}$. We show that for each system, we give its symmetry and holomorphy conditions. These symmetries, holomorphy conditions and invariant divisors are new.

1. Introduction

In [17], we presented a 4-parameter family of coupled Painlevé VI systems in dimension four with affine Weyl group symmetry of type $E_6^{(2)}$.

Now, we consider the following problem.

Problem For each affine root system $X_i^{(2)}$ with affine Weyl group $W(X_i^{(2)})$, find a system of differential equations for which $W(X_i^{(2)})$ acts as its Bäcklund transformations.

We will complete the study of the above problem in a series of papers, for which this paper is the second, resulting in a series of equations for the remaining affine root systems of types $A_l^{(2)}$ and $D_4^{(3)}$. This paper is the stage in this project where we find a 4-parameter family of coupled Painlevé VI systems in dimension four with $W(A_7^{(2)})$-symmetry explicitly given by

$$\frac{dx}{dt}=\frac{\partial H}{\partial y}, \quad \frac{dy}{dt}=-\frac{\partial H}{\partial x}, \quad \frac{dz}{dt}=\frac{\partial H}{\partial w}, \quad \frac{dw}{dt}=-\frac{\partial H}{\partial z} \tag{1}$$

with the polynomial Hamiltonian

$$\begin{aligned} t(t-1)H =&x^2y^3-((t+1)x+2\alpha_2+\alpha_3+\alpha_4)xy^2\\ &+\{tx^2+(\alpha_0+2\alpha_2+\alpha_3+\alpha_4-(\alpha_0+\alpha_1)t)x\\ &+\alpha_2(\alpha_2+\alpha_3+\alpha_4)\}y+\alpha_1tx\\ &+\frac{1}{4}[-z^2w^4+2\alpha_4zw^3+((2t-1)z^2-\alpha_4^2)w^2\\ &-2\{-\alpha_0-2\alpha_2-\alpha_3-\alpha_4+(\alpha_4+1)t\}zw-t(t-1)z^2]\\ &+y((y-1)x-\alpha_2)zw. \end{aligned} \tag{2}$$

2000 *Mathematics Subjet Classification.* 34M55; 34M45; 58F05; 32S65.

Key words and phrases. Affine Weyl group, birational symmetry, coupled Painlevé system.

Here x, y, z and w denote unknown complex variables, and $\alpha_0, \alpha_1, \dots, \alpha_4$ are complex parameters satisfying the relation:

$$\alpha_0 + \alpha_1 + 2\alpha_2 + 2\alpha_3 + \alpha_4 = 1. \tag{3}$$

This is the first example which gave higher-order Painlevé equations of type $A_{2l+5}^{(2)}$.

We then give an explicit description of a confluence process from this system to a 3-parameter family of ordinary differential systems in dimension four with $W(C_3^{(1)})$-symmetry explicitly given by

$$\frac{dx}{dt} = \frac{\partial H}{\partial y}, \quad \frac{dy}{dt} = -\frac{\partial H}{\partial x}, \quad \frac{dz}{dt} = \frac{\partial H}{\partial w}, \quad \frac{dw}{dt} = -\frac{\partial H}{\partial z} \tag{4}$$

with the polynomial Hamiltonian

$$\begin{aligned} t^2 H =& x^2y^3 - (tx + 2\alpha_1 + \alpha_2 + \alpha_3)xy^2 - \{(\alpha_0 t - 1)x - \alpha_1(\alpha_1 + \alpha_2 + \alpha_3)\}y - tx \\ & - \frac{z^2w^4}{4} + \frac{\alpha_3}{2}zw^3 + \frac{1}{4}(2tz^2 - \alpha_3^2)w^2 - \frac{1}{2}\{(\alpha_3 + 1)t - 1\}zw - \frac{t^2z^2}{4} \\ & + (xy - \alpha_1)yzw. \end{aligned} \tag{5}$$

Here x, y, z and w denote unknown complex variables, and $\alpha_0, \alpha_1, \alpha_2, \alpha_3$ are complex parameters satisfying the relation:

$$\alpha_0 + 2\alpha_1 + 2\alpha_2 + \alpha_3 = 1. \tag{6}$$

Moreover, the Bäcklund transformation group for $C_3^{(1)}$ root system can be obtained from that for type $A_7^{(2)}$ by this degeneration process.

We show that by making birational and symplectic transformations with some parameter's change this system is equivalent to a 3-parameter family of coupled Painlevé V and III systems in dimension four with $W(A_5^{(2)})$-symmetry explicitly given by

$$\frac{dx}{dt} = \frac{\partial H}{\partial y}, \quad \frac{dy}{dt} = -\frac{\partial H}{\partial x}, \quad \frac{dz}{dt} = \frac{\partial H}{\partial w}, \quad \frac{dw}{dt} = -\frac{\partial H}{\partial z} \tag{7}$$

with the polynomial Hamiltonian

$$\begin{aligned} t^2 H =& - tx(x-1)y^2 + \{x^2 + ((\alpha_0 + \alpha_1)t - 1)x - \alpha_1 t\}y + (\alpha_2 + \alpha_3)x - (\alpha_0 t - 1)\alpha_1 \\ & - \frac{z^2w^4}{4} + \frac{\alpha_3}{2}zw^3 + \frac{1}{4}(2tz^2 - \alpha_3^2)w^2 - \frac{1}{2}\{(\alpha_0 + \alpha_1 + 2\alpha_2 + 2\alpha_3)t - 1\}zw - \frac{t^2z^2}{4} \\ & - xzw. \end{aligned} \tag{8}$$

Here x, y, z and w denote unknown complex variables, and $\alpha_0, \alpha_1, \alpha_2, \alpha_3$ are complex parameters satisfying the relation:

$$\alpha_0 + \alpha_1 + 2\alpha_2 + \alpha_3 = 1. \tag{9}$$

This is the second example which gave higher-order Painlevé equations of type $A_{2l+3}^{(2)}$.

For a degenerate system of $A_5^{(2)}$ system, we find a 2-parameter family of ordinary differential systems in dimension four with $W(D_4^{(3)})$-symmetry explicitly given by

$$\frac{dx}{dt}=\frac{\partial H}{\partial y}, \quad \frac{dy}{dt}=-\frac{\partial H}{\partial x}, \quad \frac{dz}{dt}=\frac{\partial H}{\partial w}, \quad \frac{dw}{dt}=-\frac{\partial H}{\partial z} \tag{10}$$

with the polynomial Hamiltonian

(11)

$$\begin{aligned}
&(4t+3)(16t^2-12t+9)H=\\
&-12tx^2y^4+12(x+2\alpha_0 t)xy^3+3\{(\alpha_0+14\alpha_1+9\alpha_2)x-4\alpha_0^2t\}y^2\\
&+\{-24tx^2-32t^2x-3\alpha_0(5\alpha_0+14\alpha_1+9\alpha_2)\}y+24x\{x+(4\alpha_0+7\alpha_1+4\alpha_2)t\}\\
&-12tz^2w^4-12\{z-2(\alpha_0+\alpha_2)t\}zw^3+3\{8t^2z^2-(\alpha_0+6\alpha_1-3\alpha_2)z-4(\alpha_0+\alpha_2)^2t\}w^2\\
&+\{12tz^2-8(7\alpha_0+8\alpha_1+7\alpha_2)t^2z+3(5-6\alpha_1-8\alpha_1^2-4\alpha_2)\}w\\
&-3\{(4t^3+1)z+(19\alpha_0+10\alpha_1+7\alpha_2)t\}z\\
&-12x\{-4txwy^3+(4xw+4txw^2-2tzw^2+zw+2t^2z)y^2\\
&+(-2xw^2-7tz-4t^2zw+zw^2+4tzw^3)y+(8tw+5)z\}+24\alpha_0^2tyw\\
&+12\alpha_1xw(2w-7y)+6\alpha_2xyw\{4t(2w-y)-3\}\\
&+6\alpha_0y\{4t(2x-z)w^2+(x-12txy+2z)w+4t^2z\}+24\alpha_0\alpha_2tyw.
\end{aligned}$$

Here x,y,z and w denote unknown complex variables, and $\alpha_0,\alpha_1,\alpha_2$ are complex parameters satisfying the relation:

$$\alpha_0+2\alpha_1+\alpha_2=1. \tag{12}$$

This is the first example which gave higher order Painlevé type systems of type $D_4^{(3)}$.

We remark that for each system we tried to seek its first integrals of polynomial type with respect to x,y,z,w. However, we can not find. Of course, each Hamiltonian is not its first integral.

The Bäcklund transformations of each system (except for $C_3^{(1)}$-system) satisfy

$$s_i(g)=g+\frac{\alpha_i}{f_i}\{f_i,g\}+\frac{1}{2!}\left(\frac{\alpha_i}{f_i}\right)^2\{f_i,\{f_i,g\}\}+\cdots \quad (g\in\mathbb{C}(t)[x,y,z,w]), \tag{13}$$

where poisson bracket $\{,\}$ satisfies the relations:

$$\{y,x\}=\{w,z\}=1, \quad \textit{the others are } 0.$$

Since these Bäcklund transformations have Lie theoretic origin, similarity reduction of a Drinfeld-Sokolov hierarchy admits such a Bäcklund symmetry.

For each system, we also give its holomorphy conditions.

These symmetries, holomorphy conditions and invariant divisors are new.

2. $A_7^{(2)}$ SYSTEM

In this section, we study a 4-parameter family of coupled Painlevé VI systems in dimension four with affine Weyl group symmetry of type $A_7^{(2)}$ given by

$$\frac{dx}{dt}=\frac{\partial H}{\partial y},\quad \frac{dy}{dt}=-\frac{\partial H}{\partial x},\quad \frac{dz}{dt}=\frac{\partial H}{\partial w},\quad \frac{dw}{dt}=-\frac{\partial H}{\partial z} \tag{14}$$

with the polynomial Hamiltonian

$$\begin{aligned} t(t-1)H =&x^2y^3-((t+1)x+2\alpha_2+\alpha_3+\alpha_4)xy^2\\ &+\{tx^2+(\alpha_0+2\alpha_2+\alpha_3+\alpha_4-(\alpha_0+\alpha_1)t)x\\ &+\alpha_2(\alpha_2+\alpha_3+\alpha_4)\}y+\alpha_1tx\\ &+\frac{1}{4}[-z^2w^4+2\alpha_4zw^3+((2t-1)z^2-\alpha_4^2)w^2\\ &-2\{-\alpha_0-2\alpha_2-\alpha_3-\alpha_4+(\alpha_4+1)t\}zw-t(t-1)z^2]\\ &+y((y-1)x-\alpha_2)zw. \end{aligned} \tag{15}$$

Here x, y, z and w denote unknown complex variables, and $\alpha_0, \alpha_1, \dots, \alpha_4$ are complex parameters satisfying the relation:

$$\alpha_0+\alpha_1+2\alpha_2+2\alpha_3+\alpha_4=1. \tag{16}$$

This is the first example which gave higher order Painlevé type systems of type $A_7^{(2)}$.

We remark that for this system we tried to seek its first integrals of polynomial type with respect to x, y, z, w. However, we can not find. Of course, the Hamiltonian H is not the first integral.

It is known that the Painlevé VI system admits the affine Weyl group symmetry of type $B_4^{(1)}$ as the group of its Bäcklund transformations in addition to the diagram automorphisms of type $D_4^{(1)}$. The diagram automorphisms change the time variable t. However, the system (14) admits the affine Weyl group symmetry of type $A_7^{(2)}$ as the group of its Bäcklund transformations, whose generators $s_0, s_1, \dots, s_4$ are determined by the invariant divisors (2.2). Of course, these transformations do not change the time variable t.

We also remark that the iniariant divisors of the system (14) are different from the ones of a 4-parameter family of 2-coupled Painlevé VI systems in dimension four with affine Weyl group symmetry of type $E_6^{(2)}$ given in the paper [17].

Now, we show that each principal part of this Hamiltonian can be transformed into canonical Painlevé VI Hamiltonian by birational and symplectic transformations.

At first, we study the Hamiltonian system

$$\frac{dx}{dt}=\frac{\partial K_1}{\partial y},\quad \frac{dy}{dt}=-\frac{\partial K_1}{\partial x} \tag{17}$$

with the polynomial Hamiltonian

$$\begin{aligned}t(1-t)K_1 =&x^2y^3-((t+1)x+2\alpha_2+\alpha_3+\alpha_4)xy^2\\&+\{tx^2+(\alpha_0+2\alpha_2+\alpha_3+\alpha_4-(\alpha_0+\alpha_1)t)x\\&+\alpha_2(\alpha_2+\alpha_3+\alpha_4)\}y+\alpha_1 tx,\end{aligned} \tag{18}$$

where setting $z=w=0$ in the Hamiltonian H, we obtain K_1.

We transform the Hamiltonian (17) into the Painlevé VI Hamiltonian:

$$\begin{aligned}&H_{VI}(x,y,t;\beta_0,\beta_1,\beta_2,\beta_3,\beta_4)\\&=\frac{1}{t(t-1)}[y^2(x-t)(x-1)x-\{(\beta_0-1)(x-1)x+\beta_3(x-t)x\\&+\beta_4(x-t)(x-1)\}y+\beta_2(\beta_1+\beta_2)x] \quad (\beta_0+\beta_1+2\beta_2+\beta_3+\beta_4=1).\end{aligned} \tag{19}$$

Step 1: We make the change of variables:

$$x_1=y, \quad y_1=-x. \tag{20}$$

Then, we can obtain the Painlevé VI Hamiltonian:

$$H_{VI}(x_1,y_1,t;\alpha_3,\alpha_3+\alpha_4,\alpha_2,\alpha_0,\alpha_1). \tag{21}$$

Of course, the parameters α_i and β_j satisfy the relations:

$$\beta_0+\beta_1+2\beta_2+\beta_3+\beta_4=\alpha_0+\alpha_1+2\alpha_2+2\alpha_3+\alpha_4=1. \tag{22}$$

We remark that all transformations are symplectic.

Next, we study the Hamiltonian system

$$\frac{dz}{dt}=\frac{\partial K_2}{\partial w}, \quad \frac{dw}{dt}=-\frac{\partial K_2}{\partial z} \tag{23}$$

with the polynomial Hamiltonian

$$\begin{aligned}t(1-t)K_2 =&\frac{1}{4}[-z^2w^4+2\alpha_4 zw^3+((2t-1)z^2-\alpha_4^2)w^2\\&-2\{-\alpha_0-2\alpha_2-\alpha_3-\alpha_4+(\alpha_4+1)t\}zw-t(t-1)z^2],\end{aligned} \tag{24}$$

where setting $x=y=0$ in the Hamiltonian H, we obtain K_2.

Let us transform the Hamiltonian (24) into the Painlevé VI Hamiltonian.

Step 1: We make the change of variables:

$$z_1=2\sqrt{t-1}z, \quad w_1=\frac{1}{2\sqrt{t-1}}w+\frac{1}{2}. \tag{25}$$

By this transformation, in the coordinate system $(Z_1,W_1)=(1/z_1,w_1)$ two of four accessible singular points are transformed into $W_1=0$ and $W_1=1$.

Step 2: We make the change of variables:

$$z_2=-(z_1w_1-\alpha_4)w_1, \quad w_2=\frac{1}{w_1}. \tag{26}$$



FIGURE 1. This figure denotes Dynkin diagram of type $A_7^{(2)}$. The symbol in each circle denotes the invariant divisors of the system (14) (see Theorem 2.1).

Step 3: We make the change of variables:

$$\begin{aligned} &z_3 = \frac{z_2}{2t-1+2\sqrt{t(t-1)}}, \quad w_3 = (2t-1+2\sqrt{t(t-1)})w_2 + 2(1-t-\sqrt{t(t-1)}), \\ &t = \frac{2T(T-1)-(2T-1)\sqrt{T(T-1)}}{4T(T-1)}. \end{aligned} \tag{27}$$

By this transformation, in the coordinate system $(Z_2, W_2) = (1/z_3, w_3)$ the others are transformed into $W_2 = 0$ and $W_2 = \frac{1}{T}$. We remark that it is not $W_2 = \infty$ but $W_2 = 0$ because we consider in the coordinate system (z_2, w_2).

Step 4: We make the change of variables:

$$z_4 = -(z_3 w_3 - \alpha_4) w_3, \quad w_4 = \frac{1}{w_3}. \tag{28}$$

Step 5: We make the change of variables:

$$z_5 = w_4, \quad w_5 = -z_4. \tag{29}$$

Then, we can obtain the Painlevé VI Hamiltonian:

$$-\frac{1}{2} H_{VI}(z_5, w_5, T; \alpha_1+\alpha_3-1, \alpha_1+\alpha_3, \alpha_4, \alpha_0+2\alpha_2+\alpha_3, \alpha_0+2\alpha_2+\alpha_3). \tag{30}$$

Of course, the parameters α_i and β_j satisfy the relations:

$$\beta_0+\beta_1+2\beta_2+\beta_3+\beta_4 = 2(\alpha_0+\alpha_1+2\alpha_2+2\alpha_3+\alpha_4)-1 = 1. \tag{31}$$

We remark that all transformations are symplectic.

THEOREM 2.1. *The system* (14) *admits extended affine Weyl group symmetry of type* $A_7^{(2)}$ *as the group of its Bäcklund transformations, whose generators* $s_0, s_1, \dots, s_4, \pi$ *defined* *as follows*: *with the notation* $(*) := (x, y, z, w, t; \alpha_0, \alpha_1, \dots, \alpha_4)$:

(32)

$$\begin{aligned}
&s_0 : (*) \rightarrow \left(x + \frac{\alpha_0}{y-1}, y, z, w, t; -\alpha_0, \alpha_1, \alpha_2 + \alpha_0, \alpha_3, \alpha_4\right),\\
&s_1 : (*) \rightarrow \left(x + \frac{\alpha_1}{y}, y, z, w, t; \alpha_0, -\alpha_1, \alpha_2 + \alpha_1, \alpha_3, \alpha_4\right),\\
&s_2 : (*) \rightarrow \left(x, y - \frac{\alpha_2}{x}, z, w, t; \alpha_0 + \alpha_2, \alpha_1 + \alpha_2, -\alpha_2, \alpha_3 + \alpha_2, \alpha_4\right),\\
&s_3 : (*) \rightarrow \left(x + \frac{\alpha_3}{y + w^2 - t}, y, z + \frac{2\alpha_3 w}{y + w^2 - t}, w, t; \alpha_0, \alpha_1, \alpha_2 + \alpha_3, -\alpha_3, \alpha_4 + 2\alpha_3\right),\\
&s_4 : (*) \rightarrow \left(x, y, z, w - \frac{\alpha_4}{z}, t; \alpha_0, \alpha_1, \alpha_2, \alpha_3 + \alpha_4, -\alpha_4\right),\\
&\pi : (*) \rightarrow \left(-x, 1 - y, \sqrt{-1}z, \frac{w}{\sqrt{-1}}, 1 - t; \alpha_1, \alpha_0, \alpha_2, \alpha_3, \alpha_4\right).
\end{aligned}$$

We note that the Bäcklund transformations of this system satisfy

$$s_i(g) = g + \frac{\alpha_i}{f_i}\{f_i, g\} + \frac{1}{2!}\left(\frac{\alpha_i}{f_i}\right)^2 \{f_i, \{f_i, g\}\} + \cdots \quad (g \in \mathbb{C}(t)[x, y, z, w]), \tag{33}$$

where poisson bracket $\{,\}$ satisfies the relations:

$$\{y, x\} = \{w, z\} = 1, \quad \textit{the others are } 0.$$

Since these Bäcklund transformations have Lie theoretic origin, similarity reduction of a Drinfeld-Sokolov hierarchy admits such a Bäcklund symmetry.

Proposition 2.2. *This system has the following invariant divisors*:

parameter's relation	f_i
$\alpha_0 = 0$	$f_0 := y - 1$
$\alpha_1 = 0$	$f_1 := y$
$\alpha_2 = 0$	$f_2 := x$
$\alpha_3 = 0$	$f_3 := y + w^2 - t$
$\alpha_4 = 0$	$f_4 := z$

We note that when $\alpha_1 = 0$, we see that the system (14) admits a particular solution $y = 0$, and when $\alpha_3 = 0$, after we make the birational and symplectic transformations:

$$x_3 = x,\ y_3 = y + w^2 - t,\ z_3 = z - 2xw,\ w_3 = w \tag{34}$$

we see that the system (14) admits a particular solution $y_3 = 0$.

Proposition 2.3. *Let us define the following translation operators*:

$$T_1 := \pi s_0 s_2 s_3 s_4 s_3 s_2 s_0, \quad T_2 := s_0 T_1 s_0, \quad T_3 := s_2 T_2 s_2, \quad T_4 := s_4 s_3 T_3 s_3 s_4. \tag{35}$$

These translation operators act on parameters α_i *as follows*:

$$\begin{aligned}T_1(\alpha_0,\alpha_1,\dots,\alpha_4)=&(\alpha_0,\alpha_1,\dots,\alpha_4)+(-1,1,0,0,0),\\ T_2(\alpha_0,\alpha_1,\dots,\alpha_4)=&(\alpha_0,\alpha_1,\dots,\alpha_4)+(1,1,-1,0,0),\\ T_3(\alpha_0,\alpha_1,\dots,\alpha_4)=&(\alpha_0,\alpha_1,\dots,\alpha_4)+(0,0,1,-1,0),\\ T_4(\alpha_0,\alpha_1,\dots,\alpha_4)=&(\alpha_0,\alpha_1,\dots,\alpha_4)+(0,0,0,-1,2).\end{aligned} \tag{36}$$

Theorem 2.4. *Let us consider a polynomial Hamiltonian system with Hamiltonian* $K \in \mathbb{C}(t)[x,y,z,w]$. *We assume that*

($A1$) *deg*$(K)=6$ *with respect to* x,y,z,w.

($A2$) *This system becomes again a polynomial Hamiltonian system in each coordinate system* r_i $(i=0,1,\dots,4)$:

$$\begin{aligned}r_0:&x_0=\frac{1}{x},\ y_0=-((y-1)x+\alpha_0)x,\ z_0=z,\ w_0=w,\\ r_1:&x_1=\frac{1}{x},\ y_1=-(yx+\alpha_1)x,\ z_1=z,\ w_1=w,\\ r_2:&x_2=-(xy-\alpha_2)y,\ y_2=\frac{1}{y},\ z_2=z,\ w_2=w,\\ r_3:&x_3=\frac{1}{x},\ y_3=-\left((y+w^2-t)x+\alpha_3\right)x,\ z_3=z-2xw,\ w_3=w,\\ r_4:&x_4=x,\ y_4=y,\ z_4=-(zw-\alpha_4)w,\ w_4=\frac{1}{w}\end{aligned} \tag{37}$$

Then such a system coincides with the system (14) *with the polynomial Hamiltonian* (15).

By this theorem, we can also recover the parameter's relation (16).

We note that the condition ($A2$) should be read that

$$r_j(K) \quad (j=0,1,2,4), \quad r_3(K+x)$$

are polynomials with respect to x,y,z,w.

3. $C_3^{(1)}$ SYSTEM

In this section, we study a 3-parameter family of ordinary differential systems in dimension four with affine Weyl group symmetry of type $C_3^{(1)}$. At first, we consider an explicit description of a confluence process from the system (14).

Theorem 3.1. *For the system* (14) *of type* $A_7^{(2)}$, *we make the change of parameters and variables*

$$\alpha_0=\frac{1}{\varepsilon}+A_0, \quad \alpha_1=-\frac{1}{\varepsilon}, \quad \alpha_2=A_1, \quad \alpha_3=A_2, \quad \alpha_4=A_3, \tag{38}$$

$$t=\frac{T}{\varepsilon}, \quad x=\varepsilon X, \quad y=\frac{Y}{\varepsilon}, \quad z=\sqrt{\varepsilon}Z, \quad w=\frac{W}{\sqrt{\varepsilon}} \tag{39}$$

from $\alpha_0,\alpha_1,\alpha_2,\alpha_3,x,y,z,w$ to A_0,A_1,A_2,A_3,X,Y,Z,W. Then the system (14) *can also be written in the new variables X,Y,Z,W and parameters A_0,A_1,A_2,A_3 as a Hamiltonian system. This new system tends to*

$$\frac{dx}{dt}=\frac{\partial H}{\partial y},\quad \frac{dy}{dt}=-\frac{\partial H}{\partial x},\quad \frac{dz}{dt}=\frac{\partial H}{\partial w},\quad \frac{dw}{dt}=-\frac{\partial H}{\partial z} \tag{40}$$

with the polynomial Hamiltonian

$$\begin{aligned} t^2H =&x^2y^3-(tx+2\alpha_1+\alpha_2+\alpha_3)xy^2-\{(\alpha_0t-1)x-\alpha_1(\alpha_1+\alpha_2+\alpha_3)\}y-tx\\ &-\frac{z^2w^4}{4}+\frac{\alpha_3}{2}zw^3+\frac{1}{4}(2tz^2-\alpha_3^2)w^2-\frac{1}{2}\{(\alpha_3+1)t-1\}zw-\frac{t^2z^2}{4}\\ &+(xy-\alpha_1)yzw, \end{aligned} \tag{41}$$

where, for notational convenience, we have renamed A_i,X,Y,Z,W to α_i,x,y,z,w (which are not the same as the previous α_i,x,y,z,w).

Here x,y,z and w denote unknown complex variables, and $\alpha_0,\alpha_1,\alpha_2,\alpha_3$ are complex parameters satisfying the relation:

$$\alpha_0+2\alpha_1+2\alpha_2+\alpha_3=1. \tag{42}$$

We remark that for this system we tried to seek its first integrals of polynomial type with respect to x,y,z,w. However, we can not find. Of course, the Hamiltonian H is not the first integral.

Theorem 3.2. *The system* (40) *admits the affine Weyl group symmetry of type $C_3^{(1)}$ as the group of its Bäcklund transformations, whose generators S_0,S_1,S_2,S_3 defined as follows*: *with the notation $(*):=(x,y,z,w,t;\alpha_0,\alpha_1,\alpha_2,\alpha_3)$*:

$$\begin{aligned} &S_0:(*)\rightarrow\left(-\left(x+\frac{\alpha_0}{y}+\frac{1}{y^2}\right),-y,\sqrt{-1}z,\frac{w}{\sqrt{-1}},-t;-\alpha_0,\alpha_1+\alpha_0,\alpha_2,\alpha_3\right),\\ &S_1:(*)\rightarrow\left(x,y-\frac{\alpha_1}{x},z,w,t;\alpha_0+2\alpha_1,-\alpha_1,\alpha_2+\alpha_1,\alpha_3\right),\\ &S_2:(*)\rightarrow\left(x+\frac{\alpha_2}{y+w^2-t},y,z+\frac{2\alpha_2w}{y+w^2-t},w,t;\alpha_0,\alpha_1+\alpha_2,-\alpha_2,\alpha_3+2\alpha_2\right),\\ &S_3:(*)\rightarrow\left(x,y,z,w-\frac{\alpha_3}{z},t;\alpha_0,\alpha_1,\alpha_2+\alpha_3,-\alpha_3\right). \end{aligned} \tag{43}$$

We remark that the transformations S_i $(i=1,2,3)$ satisfy the relation (33) (see Section 2). However, the transformation S_0 does not satisfy so.

Theorem 3.3. *For the transformations* (38), (39) *given in Theorem* 3.1, we can choose a subgroup $W_{A_7^{(2)}\rightarrow C_3^{(1)}}$ of the Bäcklund transformation group $W(A_7^{(2)})$ so that $W_{A_7^{(2)}\rightarrow C_3^{(1)}}$ converges to the Bäcklund transformation group $W(C_3^{(1)})$ of the system (40).

Proof. Notice that

$$\alpha_0+\alpha_1+2\alpha_2+2\alpha_3+\alpha_4=A_0+2A_1+2A_2+A_3=1$$

and the change of variables from (x,y,z,w) to (X,Y,Z,W) is symplectic. Choose S_i $(i=0,1,2,3)$ as

$$S_0:=s_0s_1,\ S_1:=s_2,\ S_2:=s_3,\ S_3:=s_4,$$

where s_i are given by Theorem 2.1. Then the transformations S_i are reflections of the parameters A_0,A_1,A_2,A_3. The transformation group $W_{A_7^{(2)}\to C_3^{(1)}}=<S_0,S_1,S_2,S_3>$ coincides with the group given in Theorem 3.2 as $\varepsilon\to 0$. □

Proposition 3.4. *This system has the following invariant divisors*:

parameter's relation	f_i
$\alpha_1=0$	$f_1:=x$
$\alpha_2=0$	$f_2:=y+w^2-t$
$\alpha_3=0$	$f_3:=z$

Theorem 3.5. *Let us consider a polynomial Hamiltonian system with Hamiltonian* $K\in\mathbb{C}(t)[x,y,z,w]$. *We assume that*

$(G1)$ *deg*$(K)=6$ *with respect to* x,y,z,w.

$(G2)$ *This system becomes again a polynomial Hamiltonian system in each coordinate system* r_i $(i=0,1,3)$:

$$\begin{aligned}
&r_0:x_0=x+\frac{\alpha_0}{y}+\frac{1}{y^2},\ y_0=y,\ z_0=z,\ w_0=w,\\
&r_1:x_1=-(xy-\alpha_1)y,\ y_1=\frac{1}{y},\ z_1=z,\ w_1=w,\\
&r_2:x_2=\frac{1}{x},\ y_2=-\left((y+w^2-t)x+\alpha_2\right)x,\ z_2=z-2xw,\ w_2=w,\\
&r_3:x_3=x,\ y_3=y,\ z_3=-(zw-\alpha_3)w,\ w_3=\frac{1}{w}.
\end{aligned}\tag{44}$$

Then such a system coincides with the system (40) *with the polynomial Hamiltonian* (41).

By this theorem, we can also recover the parameter's relation (42).

We note that the conditions $(G2)$ should be read that

$$r_j(K)\quad (j=0,1,3),\quad r_2(K+x)$$

are polynomials with respect to x,y,z,w.

4. $A_5^{(2)}$ SYSTEM

In this section, we study a 3-parameter family of coupled Painlevé V and III systems in dimension four with affine Weyl group symmetry of type $A_5^{(2)}$. At first, by making birational and symplectic transformations with some parameter's change the system (40) is equivalent to the system of type $A_5^{(2)}$.



FIGURE 2. This figure denotes Dynkin diagram of type $A_5^{(2)}$. The symbol in each circle denotes the invariant divisors of the system (47) (see Theorem 4.2).

THEOREM 4.1. *For the system* (40) *of type* $C_3^{(1)}$, *we make the change of parameters and variables*

$$\alpha_0=A_0-A_1,\quad \alpha_1=A_1,\quad \alpha_2=A_2,\quad \alpha_3=A_3, \tag{45}$$

$$x=\frac{1}{X},\quad y=-(XY+A_1)X,\quad z=Z,\quad w=W \tag{46}$$

from $\alpha_0,\alpha_1,\alpha_2,\alpha_3,x,y,z,w$ *to* A_0,A_1,A_2,A_3,X,Y,Z,W. *Then the system* (40) *can also be written in the new variables* X,Y,Z,W *and parameters* A_0,A_1,A_2,A_3 *as the polynomial Hamiltonian system given by*

$$\frac{dx}{dt}=\frac{\partial H}{\partial y},\quad \frac{dy}{dt}=-\frac{\partial H}{\partial x},\quad \frac{dz}{dt}=\frac{\partial H}{\partial w},\quad \frac{dw}{dt}=-\frac{\partial H}{\partial z} \tag{47}$$

with the polynomial Hamiltonian

(48)

$$\begin{aligned}t^2H=&-tx(x-1)y^2+\{x^2+((\alpha_0+\alpha_1)t-1)x-\alpha_1t\}y+(\alpha_2+\alpha_3)x-(\alpha_0t-1)\alpha_1\\&-\frac{z^2w^4}{4}+\frac{\alpha_3}{2}zw^3+\frac{1}{4}(2tz^2-\alpha_3^2)w^2-\frac{1}{2}\{(\alpha_0+\alpha_1+2\alpha_2+2\alpha_3)t-1\}zw-\frac{t^2z^2}{4}\\&-xzw,\end{aligned}$$

where, for notational convenience, we have renamed A_i,X,Y,Z,W *to* α_i,x,y,z,w *(which are not the same as the previous* α_i,x,y,z,w*).*

Here x,y,z and w denote unknown complex variables, and $\alpha_0,\alpha_1,\alpha_2,\alpha_3$ are complex parameters satisfying the relation:

$$\alpha_0+\alpha_1+2\alpha_2+\alpha_3=1. \tag{49}$$

This is the second example which gave higher order Painlevé type systems of type $A_5^{(2)}$ (see [19]). In this case the iniariant divisors are different from the ones of a 3-parameter family of 2-coupled Painlevé III systems in dimension four given in the paper [19].

Invariant divisors	f_0	f_1	f_2	f_3
System (47)	$x-1$	x	yw^2-ty+1	z
2-CPIII	y	w	$xz+t$	$y+w-1$

We remark that for this system we tried to seek its first integrals of polynomial type with respect to x,y,z,w. However, we can not find. Of course, the Hamiltonian H is not the first integral.

We will show that each principal part of this Hamiltonian can be transformed into canonical Painlevé V and III Hamiltonian by birational and symplectic transformations. At first, we study the Hamiltonian system

$$\frac{dx}{dt}=\frac{\partial K_3}{\partial y}, \quad \frac{dy}{dt}=-\frac{\partial K_3}{\partial x} \tag{50}$$

with the polynomial Hamiltonian

(51)
$$t^2K_3=-tx(x-1)y^2+\{x^2+((\alpha_0+\alpha_1)t-1)x-\alpha_1 t\}y+(\alpha_2+\alpha_3)x-(\alpha_0 t-1)\alpha_1,$$

where setting $z=w=0$ in the Hamiltonian H, we obtain K_3.

We transform the Hamiltonian (50) into the Painlevé V Hamiltonian (see [18]):

$$\begin{aligned} &-tH_V(x,y,t;\beta_1,\beta_2,\beta_3)\\ &=-ty+x^3y^2-x^2y^2+(\beta_1+2\beta_2)x^2y+(t-1+2\beta_3)xy+\beta_2(\beta_1+\beta_2)x. \end{aligned} \tag{52}$$

Step 1: We make the change of variables:

$$x_1=\frac{y}{t}, \quad y_1=-tx, \quad T=-\frac{1}{t}. \tag{53}$$

Then, we can obtain the Painlevé V Hamiltonian:

$$H_V\left(x_1,y_1,T;\alpha_2+\alpha_3,\alpha_1,\frac{\alpha_0}{2}\right). \tag{54}$$

We remark that all transformations are symplectic.

Next, we study the Hamiltonian system

$$\frac{dz}{dt}=\frac{\partial K_4}{\partial w}, \quad \frac{dw}{dt}=-\frac{\partial K_4}{\partial z} \tag{55}$$

with the polynomial Hamiltonian

(56)
$$t^2K_4=-\frac{z^2w^4}{4}+\frac{\alpha_3}{2}zw^3+\frac{1}{4}(2tz^2-\alpha_3^2)w^2-\frac{1}{2}\{(\alpha_0+\alpha_1+2\alpha_2+2\alpha_3)t-1\}zw-\frac{t^2z^2}{4},$$

where setting $x=y=0$ in the Hamiltonian H, we obtain K_4.

Let us transform the Hamiltonian (55) into the Painlevé III Hamiltonian:

$$H_{III}(q,p,t;\alpha_1,\beta_1)=\frac{q^2p^2-(q^2-(\alpha_1+\beta_1)q-t)p-\alpha_1 q}{t}. \tag{57}$$

Step 0: We make the change of variables:

$$t=\frac{1}{T_1}. \tag{58}$$

We note that

$$dK_4\wedge dt=-\frac{1}{{T_1}^2}d\tilde{K}_4\wedge dT_1. \tag{59}$$

Step 1: We make the change of variables:

$$z_1=\frac{z}{2\sqrt{T_1}}, \quad w_1=2\sqrt{T_1}w+2. \tag{60}$$

Step 2: We make the change of variables:

$$z_2=-(z_1w_1-\alpha_3)w_1, \quad w_2=\frac{1}{w_1}. \tag{61}$$

Step 3: We make the change of variables:

$$z_3=-\frac{1}{T_1}z_2+1, \quad w_3=-T_1w_2. \tag{62}$$

Step 4: We make the change of variables:

$$z_4=w_3+\frac{T_1}{4}, \quad w_4=-z_3, \quad T_1=4T_2. \tag{63}$$

Step 5: We make the change of variables:

$$z_5=z_4, \quad w_5=w_4+1, \quad T_2=\sqrt{T_3}. \tag{64}$$

Then, we can obtain the Painlevé III Hamiltonian:

$$\frac{1}{2}H_{III}(z_5,w_5,T_3;0,\alpha_3+1). \tag{65}$$

We remark that all transformations are symplectic.

THEOREM 4.2. *The system* (47) *admits extended affine Weyl group symmetry of type* $A_5^{(2)}$ *as the group of its Bäcklund transformations, whose generators* s_0,s_1,s_2,s_3,π *defined* *as follows*: *with the notation* $(*) := (x,y,z,w,t;\alpha_0,\alpha_1,\alpha_2,\alpha_3)$:

(66)

$$\begin{aligned}
&s_0 : (*) \to \left(x, y - \frac{\alpha_0}{x-1}, z, w, t; -\alpha_0, \alpha_1, \alpha_2 + \alpha_0, \alpha_3\right),\\
&s_1 : (*) \to \left(x, y - \frac{\alpha_1}{x}, z, w, t; \alpha_0, -\alpha_1, \alpha_2 + \alpha_1, \alpha_3\right),\\
&s_2 : (*) \to \left(x + \frac{\alpha_2(w^2-t)}{yw^2-ty+1}, y, z + \frac{2\alpha_2 yw}{yw^2-ty+1}, w, t; \alpha_0 + \alpha_2, \alpha_1 + \alpha_2, -\alpha_2, \alpha_3 + 2\alpha_2\right),\\
&s_3 : (*) \to \left(x, y, z, w - \frac{\alpha_3}{z}, t; \alpha_0, \alpha_1, \alpha_2 + \alpha_3, -\alpha_3\right),\\
&\pi : (*) \to \left(1-x, -y, \sqrt{-1}z, \frac{w}{\sqrt{-1}}, -t; \alpha_1, \alpha_0, \alpha_2, \alpha_3\right).
\end{aligned}$$

These Bäcklund transformations have Lie theoretic origin, similarity reduction of a Drinfeld-Sokolov hierarchy admits such a Bäcklund symmetry.

Proposition 4.3. *This system has the following invariant divisors*:

parameter's relation	f_i
$\alpha_0 = 0$	$f_0 := x - 1$
$\alpha_1 = 0$	$f_1 := x$
$\alpha_2 = 0$	$f_2 := yw^2 - ty + 1$
$\alpha_3 = 0$	$f_3 := z$

We note that the system (47) admits a Riccati extension of the fifth Painlevé system as its particular solutions when $z = 0$ with the parameter's relation $\alpha_3 = 0$.

Proposition 4.4. *Let us define the following translation operators*:

$$T_1 := \pi s_0 s_2 s_3 s_2 s_0, \quad T_2 := \pi s_0 s_1 s_2 s_3 s_2, \quad T_3 := s_2 T_2 s_2. \tag{67}$$

These translation operators act on parameters α_i *as follows*:

$$\begin{aligned}
T_1(\alpha_0,\alpha_1,\alpha_2,\alpha_3) =&(\alpha_0,\alpha_1,\alpha_2,\alpha_3) + (-1,1,0,0),\\
T_2(\alpha_0,\alpha_1,\alpha_2,\alpha_3) =&(\alpha_0,\alpha_1,\alpha_2,\alpha_3) + (1,1,-1,0),\\
T_3(\alpha_0,\alpha_1,\alpha_2,\alpha_3) =&(\alpha_0,\alpha_1,\alpha_2,\alpha_3) + (0,0,1,-2).
\end{aligned} \tag{68}$$

Theorem 4.5. *Let us consider a polynomial Hamiltonian system with Hamiltonian* $K \in \mathbb{C}(t)[x,y,z,w]$. *We assume that*

$(B1)$ $deg(K) = 6$ *with respect to* x, y, z, w.

($B2$) *This system becomes again a polynomial Hamiltonian system in each coordinate system* r_i $(i=0,1,3)$:

$$\begin{aligned} &r_0 : x_0 = -((x-1)y-\alpha_0)y,\ y_0 = \frac{1}{y},\ z_0 = z,\ w_0 = w,\\ &r_1 : x_1 = -(xy-\alpha_1)y,\ y_1 = \frac{1}{y},\ z_1 = z,\ w_1 = w,\\ &r_3 : x_3 = x,\ y_3 = y,\ z_3 = -(zw-\alpha_3)w,\ w_3 = \frac{1}{w}. \end{aligned} \tag{69}$$

($B3$) *In addition to the assumption* ($B2$), *the Hamiltonian system in the coordinate* r_1 *becomes again a polynomial Hamiltonian system in the coordinate system* r_2:

$$r_2 : x_2 = \frac{1}{x_1},\ y_2 = -\left((y_1 + w_1^2 - t)x_1 + \alpha_2\right)x_1,\ z_2 = z_1 - 2x_1 w_1,\ w_2 = w_1.$$

Then such a system coincides with the system (47) *with the polynomial Hamiltonian* (48).

By this theorem, we can also recover the parameter's relation (49).

We note that the conditions ($B2$) and ($B3$) should be read that

$$r_j(K) \quad (j=0,1,3), \quad r_2(r_1(K)+x_1)$$

are polynomials with respect to x,y,z,w or x_1,y_1,z_1,w_1.

5. $D_4^{(3)}$ SYSTEM

In this section, we study a 2-parameter family of ordinary differential systems in dimension four with affine Weyl group symmetry of type $D_4^{(3)}$ given by

$$\frac{dx}{dt} = \frac{\partial H}{\partial y}, \quad \frac{dy}{dt} = -\frac{\partial H}{\partial x}, \quad \frac{dz}{dt} = \frac{\partial H}{\partial w}, \quad \frac{dw}{dt} = -\frac{\partial H}{\partial z} \tag{70}$$

with the polynomial Hamiltonian

(71)

$$\begin{aligned} &(4t+3)(16t^2-12t+9)H =\\ &-12tx^2y^4 + 12(x+2\alpha_0 t)xy^3 + 3\{(\alpha_0+14\alpha_1+9\alpha_2)x - 4\alpha_0^2 t\}y^2\\ &+\{-24tx^2 - 32t^2x - 3\alpha_0(5\alpha_0+14\alpha_1+9\alpha_2)\}y + 24x\{x+(4\alpha_0+7\alpha_1+4\alpha_2)t\}\\ &-12tz^2w^4 - 12\{z-2(\alpha_0+\alpha_2)t\}zw^3 + 3\{8t^2z^2 - (\alpha_0+6\alpha_1-3\alpha_2)z - 4(\alpha_0+\alpha_2)^2t\}w^2\\ &+\{12tz^2 - 8(7\alpha_0+8\alpha_1+7\alpha_2)t^2z + 3(5-6\alpha_1-8\alpha_1^2-4\alpha_2)\}w\\ &-3\{(4t^3+1)z + (19\alpha_0+10\alpha_1+7\alpha_2)t\}z\\ &-12x\{-4txwy^3 + (4xw+4txw^2-2tzw^2+zw+2t^2z)y^2\\ &+(-2xw^2-7tz-4t^2zw+zw^2+4tzw^3)y + (8tw+5)z\} + 24\alpha_0^2tyw\\ &+12\alpha_1xw(2w-7y) + 6\alpha_2xyw\{4t(2w-y)-3\}\\ &+6\alpha_0y\{4t(2x-z)w^2 + (x-12txy+2z)w + 4t^2z\} + 24\alpha_0\alpha_2tyw. \end{aligned}$$

Here x, y, z and w denote unknown complex variables, and $\alpha_0, \alpha_1, \alpha_2$ are complex parameters satisfying the relation:

$$\alpha_0 + 2\alpha_1 + \alpha_2 = 1. \tag{72}$$

This is the first example which gave higher order Painlevé type systems of type $D_4^{(3)}$.

We remark that for this system we tried to seek its first integrals of polynomial type with respect to x, y, z, w. However, we can not find. Of course, the Hamiltonian H is not the first integral.

This system can be obtained by connecting the invariant divisors $x - 1$ and z for the canonical variables (x, y, z, w).

It is still an open question whether we make an explicit description of a confluence process from the system (47) to this system.



FIGURE 3. This figure denotes Dynkin diagram of type $D_4^{(3)}$. The symbol in each circle denotes the invariant divisors of the system (70) (see Theorem 5.1).

THEOREM 5.1. *The system* (70) *admits the affine Weyl group symmetry of type* $D_4^{(3)}$ *as the group of its Bäcklund transformations, whose generators* s_0, s_1, s_2 *defined as follows*: *with the notation* $(*) := (x, y, z, w, t; \alpha_0, \alpha_1, \alpha_2)$:

$$\begin{aligned} &s_0 : (*) \rightarrow \left(x, y - \frac{\alpha_0}{x}, z, w, t; -\alpha_0, \alpha_1 + \alpha_0, \alpha_2\right), \\ &s_1 : (*) \rightarrow \left(x + \frac{\alpha_1(w^2 - t)}{yw^2 - ty + 1}, y, z + \frac{2\alpha_1 yw}{yw^2 - ty + 1}, w, t; \alpha_0 + \alpha_1, -\alpha_1, \alpha_2 + 3\alpha_1\right), \\ &s_2 : (*) \rightarrow \left(x, y - \frac{\alpha_2}{x + z}, z, w - \frac{\alpha_2}{x + z}, t; \alpha_0, \alpha_1 + \alpha_2, -\alpha_2\right). \end{aligned} \tag{73}$$

These Bäcklund transformations have Lie theoretic origin, similarity reduction of a Drinfeld-Sokolov hierarchy admits such a Bäcklund symmetry.

PROPOSITION 5.2. *This system has the following invariant divisors*:

parameter's relation	f_i
$\alpha_0 = 0$	$f_0 := x$
$\alpha_1 = 0$	$f_1 := yw^2 - ty + 1$
$\alpha_2 = 0$	$f_2 := x + z$

THEOREM 5.3. *Let us consider a polynomial Hamiltonian system with Hamiltonian* $K \in \mathbb{C}(t)[x,y,z,w]$. *We assume that*

$(C1)$ *$deg(K)=6$ with respect to x,y,z,w.*

$(C2)$ *This system becomes again a polynomial Hamiltonian system in each coordinate system r_i $(i=0,2)$:*

$$\begin{aligned} &r_0 : x_0=-(xy-\alpha_0)y,\ y_0=\frac{1}{y},\ z_0=z,\ w_0=w,\\ &r_2 : x_2=x,\ y_2=y-w,\ z_2=-((z+x)w-\alpha_2)w,\ w_2=\frac{1}{w}. \end{aligned} \tag{74}$$

$(C3)$ *In addition to the assumption $(C2)$, the Hamiltonian system in the coordinate r_1 becomes again a polynomial Hamiltonian system in the coordinate system r_0:*

$$r_1 : x_1=\frac{1}{x_0},\ y_1=-\left((y_0+w_0^2-t)x_0+\alpha_1\right)x_0,\ z_1=z_0-2x_0w_0,\ w_1=w_0.$$

Then such a system coincides with the system (70) *with the polynomial Hamiltonian* (71)*.*

By this theorem, we can also recover the parameter's relation (72).

We note that the conditions $(C2)$ and $(C3)$ should be read that

$$r_j(K) \quad (j=0,2), \quad r_2(r_0(K)+x_0)$$

are polynomials with respect to x,y,z,w or x_0,y_0,z_0,w_0.

6. APPENDIX

In this appendix, we present a 2-parameter family of ordinary differential systems with affine Weyl group symmetry of type $G_2^{(1)}$ given by

$$\frac{dx}{dt}=\frac{\partial H}{\partial y},\quad \frac{dy}{dt}=-\frac{\partial H}{\partial x},\quad \frac{dz}{dt}=\frac{\partial H}{\partial w},\quad \frac{dw}{dt}=-\frac{\partial H}{\partial z} \tag{75}$$

with the polynomial Hamiltonian

$$\begin{aligned} H=&-3(5\alpha_0+18\alpha_1+27\alpha_2)txy^3-\frac{xy}{2t}+6\alpha_1(\alpha_0+6\alpha_1+9\alpha_2)ty^2\\ &-12t^2z^2w^2-\left\{24(\alpha_1+\alpha_2)t^2-\frac{3}{2t}\right\}zw\\ &+12t^2y^3zw^2-24txy^3zw+12x^2y^3z+24txz^2w-12x^2z^2-12\alpha_0t^2y^3w\\ &+12(2\alpha_0+4\alpha_1+9\alpha_2)xy^2z+3(19\alpha_0+38\alpha_1+45\alpha_2)ty^2zw\\ &-6\alpha_1(\alpha_0+4\alpha_1+9\alpha_2)yz-4(\alpha_0-4\alpha_1-3\alpha_2)txz. \end{aligned} \tag{76}$$

Here x,y,z and w denote unknown complex variables, and $\alpha_0,\alpha_1,\alpha_2$ are complex parameters satisfying the relation:

$$\alpha_0+2\alpha_1+3\alpha_2=0. \tag{77}$$

This is the first example which gave higher order Painlevé type systems of type $G_2^{(1)}$.

We remark that for this system we tried to seek its first integrals of polynomial type with respect to x, y, z, w. However, we can not find. Of course, the Hamiltonian H and $z - ty$ are not its first integrals.



FIGURE 4. The symbol in each circle denotes the invariant divisors of this system (see Theorem 6.1).

THEOREM 6.1. *The system* (75) *admits the affine Weyl group symmetry of type* $G_2^{(1)}$ *as the group of its Bäcklund transformations, whose generators* s_0, s_1, s_2 *defined as follows: with the notation* $(*) := (x, y, z, w, t; \alpha_0, \alpha_1, \alpha_2)$:

$$\begin{aligned} &s_0 : (*) \rightarrow \left(x, y, z, w - \frac{\alpha_0}{z}, t; -\alpha_0, \alpha_1 + \alpha_0, \alpha_2\right), \\ &s_1 : (*) \rightarrow \left(x, y - \frac{\alpha_1}{x - tw}, z - \frac{\alpha_1 t}{x - tw}, w, t; \alpha_0 + \alpha_1, -\alpha_1, \alpha_2 + \alpha_1\right), \\ &s_2 : (*) \rightarrow \left(x - \frac{3\alpha_2 y^2}{z - y^3}, y, z, w - \frac{\alpha_2}{z - y^3}, t; \alpha_0, \alpha_1 + 3\alpha_2, -\alpha_2\right). \end{aligned} \tag{78}$$

Since these Bäcklund transformations have Lie theoretic origin, similarity reduction of a Drinfeld-Sokolov hierarchy admits such a Bäcklund symmetry.

PROPOSITION 6.2. *This system has the following invariant divisors*:

parameter's relation	f_i
$\alpha_0 = 0$	$f_0 := z$
$\alpha_1 = 0$	$f_1 := x - tw$
$\alpha_2 = 0$	$f_2 := z - y^3$

THEOREM 6.3. *Let us consider a polynomial Hamiltonian system with Hamiltonian* $K \in \mathbb{C}(t)[x, y, z, w]$. *We assume that*

$(F1)$ *deg*$(K) = 6$ *with respect to* x, y, z, w.

$(F2)$ *This system becomes again a polynomial Hamiltonian system in each coordinate system* r_i $(i = 0, 1, 2)$:

$$\begin{aligned} &r_0 : x_0 = x, \ y_0 = y, \ z_0 = -(zw - \alpha_0)w, \quad w_0 = \frac{1}{w}, \\ &r_1 : x_1 = -((x - tw)y - \alpha_1)y, \ y_1 = \frac{1}{y}, \ z_1 = z - ty, \ w_1 = w, \\ &r_2 : x_2 = x - 3y^2 w, \ y_2 = y, \ z_2 = -((z - y^3)w - \alpha_2)w, \ w_2 = \frac{1}{w}. \end{aligned} \tag{79}$$

Then such a system coincides with the system

$$\frac{dx}{dt}=\frac{\partial K}{\partial y}, \quad \frac{dy}{dt}=-\frac{\partial K}{\partial x}, \quad \frac{dz}{dt}=\frac{\partial K}{\partial w}, \quad \frac{dw}{dt}=-\frac{\partial K}{\partial z} \tag{80}$$

with the polynomial Hamiltonian

$$K = H + \sum_{i=1}^{6} a_i(z-ty)^i \quad (a_i \in \mathbb{C}(t)). \tag{81}$$

We note that the condition $(F2)$ should be read that

$$r_j(K) \quad (j=0,2), \quad r_1(K-yw)$$

are polynomials with respect to x, y, z, w.

REFERENCES

[1] P. Painlevé, *Mémoire sur les équations différentielles dont l'intégrale générale est uniforme*, Bull. Société Mathématique de France. **28** (1900), 201–261.

[2] P. Painlevé, *Sur les équations différentielles du second ordre et d'ordre supérieur dont l'intégrale est uniforme*, Acta Math. **25** (1902), 1–85.

[3] B. Gambier, *Sur les équations différentielles du second ordre et du premier degré dont l'intégrale générale est à points critiques fixes*, Acta Math. **33** (1910), 1–55.

[4] C. M. Cosgrove and G. Scoufis, *Painlevé classification of a class of differential equations of the second order and second degree*, Studies in Applied Mathematics. **88** (1993), 25-87.

[5] C. M. Cosgrove, *All binomial-type Painlevé equations of the second order and degree three or higher*, Studies in Applied Mathematics. **90** (1993), 119-187.

[6] F. Bureau, *Integration of some nonlinear systems of ordinary differential equations*, Annali di Matematica. **94** (1972), 345–359.

[7] J. Chazy, *Sur les équations différentielles dont l'intégrale générale est uniforme et admet des singularités essentielles mobiles*, Comptes Rendus de l'Académie des Sciences, Paris. **149** (1909), 563–565.

[8] J. Chazy, *Sur les équations différentielles dont l'intégrale générale posséde une coupure essentielle mobile* , Comptes Rendus de l'Académie des Sciences, Paris. **150** (1910), 456–458.

[9] J. Chazy, *Sur les équations différentielles du trousiéme ordre et d'ordre supérieur dont l'intégrale a ses points critiques fixes*, Acta Math. **34** (1911), 317–385.

[10] Y. Sasano, *Coupled Painlevé VI systems in dimension four with affine Weyl group symmetry of types* $B_6^{(1)}, D_6^{(1)}$ *and* $D_7^{(2)}$, preprint.

[11] Y. Sasano, *Four-dimensional Painlevé systems of types* $D_5^{(1)}$ *and* $B_4^{(1)}$, preprint.

[12] Y. Sasano, *Higher order Painlevé equations of type* ${D_l}^{(1)}$, RIMS Kokyuroku **1473** (2006), 143–163.

[13] Y. Sasano, *Symmetries in the system of type* $D_4^{(1)}$, preprint.

[14] Y. Sasano, *Coupled Painlevé III systems with affine Weyl group symmetry of types* $B_4^{(1)}$, $D_4^{(1)}$ *and* $D_5^{(2)}$, preprint.

[15] Y. Sasano, *Coupled Painlevé III systems with affine Weyl group symmetry of types* $B_5^{(1)}, D_5^{(1)}$ *and* $D_6^{(2)}$, preprint.

[16] Y. Sasano, *Coupled Painlevé VI systems in dimension four with affine Weyl group symmetry of type* $D_6^{(1)}$*, II*, RIMS Kokyuroku Bessatsu. **B5** (2008), 137–152.

[17] Y. Sasano, *Coupled Painlevé VI systems in dimension four with affine Weyl group symmetry of type* $E_6^{(2)}$, preprint.

[18] Y. Sasano, *Symmetry in the Painlevé systems and their extensions to four-dimensional systems*, Funkcial. Ekvac. **51** (2008), 351–369.

[19] Y. Sasano, *Symmetries in the system of type* $A_5^{(2)}$, preprint.